Report No.: 8002-06-4

# CONFIDENTIAL NOT FOR PUBLIC RELEASE

#### RECOMMENDATION

Based on information contained in the site inspection report and additional information developed from ongoing UST investigations, groundwater and soils exposure are the pathways of concern. Although groundwater samples were not collected during inspection or investigation activities, groundwater is a principle source of drinking water within 4 miles of site, serving a population of approximately 158,066. The nearest well is located approximately 0.6 miles east of the site. Additionally, the site is situated within a well head protection area. There are currently 100 employees on-site and the combined population in the four-mile vicinity of the site is 182,975. The above information supports a recommendation of **Lower Priority for Further Action** for the John Hassall Inc. site.

DECLASSIFIED


JA

8/29/22
Initials/Date

333202

# APPENDIX D SITE INSPECTION WORKSHEETS

John Hassall Inc. 8002-064 June 30, 1992 Site Name:



#### GENERAL INFORMATION (continued)

Source Descriptions:

POTENTIAL WASTE UNITS:

Waste Unit 1 - Surface Impoundment: Identified in SI report (Ref. No. 28 p. 2)

Waste Unit 2 - Underground Storage Tanks (UST): Identified in SI report (Ref. No. 28 p. 2)

Hazardous Waste Quantity (HWQ) Calculations:

(See Si Tables 1 and 2)

Waste Unit 1 - Surface Impoundment: This potential waste unit was incorrectly identified in the SI report. This is not an impoundment, but a recharge basin which is used to provide rapid percolation of surficial waters into the aquifer. Additionally, Surface Impoundments as defined in current PA, SI and HRS guidance indicate a storage of accumulated waste. This is opposite of the purpose of a recharge basin.

Waste Unit 2 - Underground Storage Tank (UST): USTs #5,7,8 (10,000 gallons each) and 11 (1000 gallons) have been identified as leaking (Ref. Nos. 7,8,9). The contents of the tanks has been identified as chromium, copper, ehtylbenzene, xylenes (m,p and o) and nickel contaminated waste waters from the plating processes.

Total Gallons = 31,000 HWQ = 10

wa = 10

Attach additional pages, if necessary.


## SI TABLE 1: HAZARDOUS WASTE QUANTITY (HWQ) SCORE 60 1992 SINGLE SCURCE SITES AND FORMULAS FOR MULTIPLE SOURCE SITES

|             |                                   |   | · · · · · · · · · · · · · · · · · · ·                       |  |  |   |   |
|-------------|-----------------------------------|---|---|--|--|---|---|
| T -         |                                   |   | SI  | NGLE SOURCE SITES  | (assigned HWQ scores   | .)  | MULTIPLE<br>SOURCE<br>SITES                       |
| E<br>R      | SOURCE TYPE                       | 4 | HWQ = 10  | HWQ = 100  | HWQ = 10,000   | H <b>WQ ≃</b><br>1,000,000                                      | Formula for<br>Assigning<br>Source WQ<br>Values   |
| COMSTITUENT | N/A                               |   | ವ 100 ನೀಕ   | > 100 to 10,000 the  | > 10,000 to 1,000,000 fbe  | > 1,000,000 lbe   | lbs + 1   |
| ¥457657464¥ | N/A                               |   | ≤ 500,000 fbs   | > 500,000 to 50 million (bs  | >50 million to 5 billion lbs   | > 5 billion (bs   | /bs + 5,000                                       |
|             | Langfil                           |   | ≤ 6.75 million ft <sup>3</sup><br>≤ 250,000 ya <sup>3</sup> | > 6.75 million to 675 million ft <sup>2</sup><br>> 250,000 to 25 million ya <sup>2</sup> | > 675 million to 67.5 billion tt <sup>2</sup><br>> 25 million to 2.5 billion yd <sup>3</sup> | > 67.5 billion ft <sup>3</sup><br>> 2.5 billion yd <sup>3</sup> | $ft^3 + 67,50.$<br>$yd^3 + 2,50.$                 |
| v           | Surface<br>impoundment            |   | ≤ 6.750 tt²<br>≤ 250 yd²                                    | > 6.750 to 675,000 ft <sup>2</sup><br>> 250 to 25,000 ye <sup>2</sup>                    | > 675,000 to 67.5 million ft <sup>3</sup><br>> 25,000 to 2.5 million yd <sup>3</sup>         | > 67.5 milion ft <sup>3</sup><br>> 2.5 million ye <sup>3</sup>  | ft <sup>3</sup> + 67.5<br>yd <sup>3</sup> + 2.5   |
| 0           | Drums                             |   | ≤ 1,000 drums   | >1,000 to 100,000 drums  | > 100,000 to 10 million drume  | > 10 million drums  | drums + 10  |
| N N         | Tanks and non-<br>drum containers |   | ±50,000 gallons   | >50,000 to 5 million gallons   | > 5 million to 500 million<br>gallons  | > 500 million<br>gallons  | gallons + 5C                                      |
| £           | Contaminated soil                 |   | ≤6.75 milion It <sup>3</sup><br>≤250,000 ye <sup>3</sup>    | > 6.75 million to 675 million to<br>> 250,000 to 25 million yel                          | > 675 million to 67.5 billion ft <sup>3</sup><br>> 25 million to 2.5 billion ye <sup>3</sup> | > 67.5 billion ft <sup>3</sup><br>> 2.5 billion yd <sup>3</sup> | ft <sup>3</sup> + 67.50<br>yd <sup>3</sup> + 2.50 |
|             | คข•                               |   | ≤6,750 tc²<br>≤250 yes²                                     | >6,750 to 675,000 ft <sup>3</sup><br>>250 to 25,000 ye <sup>3</sup>                      | > 675,000 to 67.5 million ft <sup>1</sup><br>> 25,000 to 2,5 million ya <sup>1</sup>         | >67.5 milion ft <sup>3</sup><br>>2.5 milion ye <sup>3</sup>     | ft <sup>2</sup> + 67.5<br>yd <sup>3</sup> + 2.5   |
|             | Lancti                            |   | ≤ 340,000 ft <sup>8</sup><br>≤ 7.8 acres                    | >340,000 to 34 million ft <sup>2</sup><br>>7.8 to 780 acres                              | > 34 million to 3,4 billion ft <sup>2</sup><br>> 780 to 78,000 acres                         | > 3,4 billion ft <sup>d</sup><br>> 78,000 acres                 | ft <sup>2</sup> + 3,400<br>acres +<br>0.078       |
|             | Surface<br>impoundment            |   | ≤ 1,300 ft <sup>2</sup><br>≤ 0.029 ecres                    | >1,300 to 130,000 ft <sup>3</sup><br>>0,029 to 2,9 acres                                 | > 130,000 to 13 million ft <sup>2</sup><br>> 2.9 to 290 scres                                | > 13 million ft <sup>2</sup><br>> 290 acres                     | ft <sup>3</sup> + 13<br>acres +<br>0.00029        |
| A<br>R<br>E | Contaminated soil                 |   | ≤3.4 milian ft <sup>i</sup><br>≤78 acres                    | > 3.4 million to 340 million ft <sup>4</sup><br>> 78 to 7,800 acres                      | > 340 million to 34 billion ft <sup>d</sup><br>> 7,800 to 780,000 acres                      | > 34 billion ft <sup>a</sup><br>> 780,000 scress                | ft + 34,00<br>acres + 0.7                         |
| ^           | Pile *                            |   | ≤ 1,300 ft <sup>2</sup><br>≤ 0,025 ecres                    | >1,300 to 130,000 ft <sup>2</sup><br>>0.029 to 2.9 acres                                 | > 130,000 to 13 million tt <sup>2</sup><br>> 2.9 to 290 ecres                                | > 13 million ft <sup>2</sup><br>> 290 acres                     | /t + 13<br>acres +<br>0.00029                     |
|             | Land treatment                    |   | ≤ 27,000 ft²<br>≤ 0.62 acres                                | >27,000 to 2.7 million ft <sup>3</sup><br>>0.62 to 62 acres                              | > 2.7 million to 270 million ft <sup>2</sup><br>> 62 to 6,200 scree                          | > 270 million ft <sup>1</sup><br>> 6,200 scree                  | fr + 270<br>scres +<br>0.0062                     |

1 ton = 2,000 bs = 1 yd2 = 4 drums = 200 gallone

#### SI TABLE 2: HWQ SCORES FOR MULTIPLE SOURCE SITES

| Site WQ Total          | HWQ Score |
|------------------------|-----------|
| >0 te 100              | :0        |
| > 100 to 10,000        | 100       |
| > 10,000 to 1 milion : | 10,000    |
| 5 th medical           | 1 559 159 |



<sup>\*</sup> Use area of land surface under pile, not surface area of pile.

SI TABLE 3: SOURCE HAZARDOUS SUBSTANCE SUMMARY 6 Source ID: UNDERGROUND STORAGE TANKS (31,000 Gallons) 11 ach additional pages if necessary. Aur Ecotomaty/ Persistance/ Topicity/ Topicity/ Toxicity/ Persistance/ Ecobiosccumulation Mobility Mobility Persistence Biosceumusuon Toxian Hazardous Substance 5x104 5× 104 NA 100 10,000 1 10,000 1 Chromium 5 X 106 NA Copper 200 2000 10 4.0 0.1 thulbenzene 10 1.0 2 x 104 1.0 | 0.01 0.4 2000 1.0 20 1.0 | 0.01 0.4 2000 1.0 0.4 1.010.01 20 NA 5×103 5×10 10,000 10,000 100 5×106 5×104 10 10,000 | 100 10,000 Highest Values Ref. No. 9 Source ID: Highest Values Source ID: Highest Values Highest Values IAM Sources

ω

Population Served

Reference

Dose

Percent of

Cancer Risk

Concentration

Cancer Risk

Cancentration

Percent of

Reference

Dase

| SI TABLE 4: GRO | OUND WATER OBSERVED          | RELEASE SU            | JBSTANCES | No GW samples to                                 |                 |
|-----------------|------------------------------|-----------------------|-----------|--|-----------------|
| Sample ID       | Hazardoua<br>Substanca       | Toxicity/<br>Mobility | Reference | No GW samples to<br>support GW<br>contamination. | VO <sub>V</sub> |
| <u></u>         | JA:                          |                       |           | (Ref Nos. 4; 24; 26)                             | <u>:33</u>      |
|                 | Highest<br>Toxicity/Mobility |                       |           |  |                 |

Level I \_\_\_\_

Benchmark

Concentration

(MCL or MCLG)

Concentration

(ug/L)

#### SI TABLE 5: GROUND WATER ACTUAL CONTAMINATION TARGETS

Hazardous

Substance

Well ID:

Sample ID


|           | VF                     |                         |   |                         |                              |  |                    |                                 |
|-----------|------------------------|-------------------------|---|-------------------------|------------------------------|--|--------------------|---------------------------------|
|           |                        |                         | flighest<br>Percent                         |                         | Sum of<br>Percents           |  | Sum of<br>Percente |                                 |
| Well ID:  |                        |                         | Level i                                     | Level II                | _                            | Po   | pulation Serve     | d                               |
| Sample ID | Hezerdous<br>Substance | Concentration<br>(ug/L) | Benchmark<br>Concentration<br>(MCL or MCLG) | Percent of<br>Benchmark | Cancer Risk<br>Concentration | Parcent of<br>Cancer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose |
|           | NA                     |                         |   |                         |                              |  |                    |                                 |
|           |                        |                         | Highest<br>Percent                          |                         | Sum of<br>Percents           |  | Sum of<br>Percents |                                 |

Level II

Percent of

Benchmark

|     | GROUND WATER PATHWAY WORKSHEET  |        |              | 9                       |
|-----|---|--------|--------------|-------------------------|
|     | nen 6 <b>1991</b>   |        | Data         |                         |
| LIK | ELIHOOD OF RELEASE  | Score  | Type         | Referen                 |
| 1.  | CBSERVED RELEASE: If sampling data or direct observation support a release to ground water, assign a score of 550. Record observed release substances on SI Table 4.  |        |              |                         |
| 2.  | NO OBSERVED RELEASE: If sampling data do not support a release to ground water, and the site is in karst terrain or the depth to aquifer is 70 feet or less, assign a score of 500; otherwise, assign a score of 340.   | 500    |              |                         |
|     | LR =  | 500    |              |                         |
| TA  | RGETS   |        | <del>,</del> | •                       |
|     | Are any wells part of a blended system? Yes No lf yes, attach a page to show apportionment calculations.  |        |              |                         |
| 3.  | ACTUAL CONTAMINATION TARGETS: If analytical evidence indicates that any drinking-<br>water well has been exposed to a hazardous substance from the site, calculate the factor<br>score based on the number of people served by using SI Table 5.  |        |              | Ref                     |
|     | Level I: people x 10 = Total =  | . 0    |              | 10                      |
| 4.  | POTENTIAL CONTAMINATION TARGETS: Determine the number of people served by drinking-water wells that are not exposed to a hazardous substance from the site; record the population for each distance category in SI Table 6a or 6b, and assign the total population score.   | 3,301  |              | Ref<br>Nos<br>13;<br>23 |
| 5.  | NEAREST WELL: Assign a score of 50 for any Level I Actual Contamination Targets.  Assign a score of 45 if there are Level II targets but no Level I targets. If no Actual Contamination Targets exist, assign the Nearest Well score from SI Table 6a or 6b. If no drinking-water wells exist within 4 miles, assign 0. | 9      |              | Ref<br>No<br>27         |
| 6.  | WELLHEAD PROTECTION AREA (WHPA): If any source lies within or above a WHPA, or if a ground water observed release has occurred within a WHPA, assign a score of 20; assign 5 if neither condition applies but a WHPA is within 4 miles; otherwise assign 0.   | 20     |              | 20f<br>120<br>83        |
| 7.  | RESOURCES   | 5      |              |                         |
| ш   | Т:  | 12,235 |              |                         |



Site Name: Date:

## SI TABLE 6: VALUES FOR POTENTIAL CONTAMINATION GROUND WATER TARGET POPULATIONS

SI Table 6a: Non-Karst Aquilers

|                |            | Nearest | <u> </u>   | Pos        | nulation | Serve | d by W | ells Withi | n Distance | Categor | Υ       |         | <del></del> |           |                     |
|----------------|------------|---------|------------|------------|----------|-------|--------|------------|------------|---------|---------|---------|-------------|-----------|---------------------|
|                |            | Well    | ,          | 77         | 31       | 101   | 301    | 1,001      | 3,001      | 10.001  | 30,001  | 100,001 | 300.001     | 1,000,001 | Omerstellen         |
| istance        |            | (choase | 10         | te         | 10       | 10    | 10     | t-         | io         | 10      | to.     | 10      | 1.000.000   | 1 000 000 | Population<br>Value |
| am Site        | Population | Nghesti | 10         | 30         | 100      | 300   | 1.000  | 3,000      | 10,000     | 30,000  | 100,000 | 300,000 | 7.000.000   | 3.000,000 |                     |
|                | 0          | 20      | 1          | 2          | 5        | 16    | 52     | 163        | 521        | 1,633   | 5,214   | 16,325  | 62,136      | 163,246   |                     |
| to % mile      |            | • •     | ١.         | ١,         | 3        | 10    | 32     | 101        | 323        | 1,012   | 3,233   | 10,121  | 32,324      | 101,212   |                     |
| to is mile     | 0          | 18      | l '        | \          |          | ] '   |        |            |            |         | 1,668   | 5,224   | 16,684      | 52,239    | 167                 |
| is to I mile   | 9,952      | 9       | 1          | ١ ١        | 2        | 6     | 17     | 62         | (167)      | 522     | 1,008   | 3,224   |             | 1         | 939                 |
|                | 31,782     | 5       | 1,         | ,          | 1        | 3     | 9      | 29         | 94         | 294     | 939     | 2,938   | 9,385       | 29,384    |                     |
| · 1 to 2 miles | 49,548     | 1       |            | ١.         | ١.       | ١,    | ١,     | 21         | 68         | 212     | 678     | 2,122   | 6,777       | 21,222    | 678                 |
| - 2 to 3 miles |            |         | <b>!</b> ' | <b>\</b> ' | ' '      | 1 1   | 1      | 1 ~        |            | ì       |         |         | 4,171       | 13,060    | 417                 |
| · 3 to 4 miles | 64,868     | 2       | ١,         | 1          | 1        | 1     | 1 4    | 13         | 42         | 131     | (11)    | 1,308   | 1 7.1/1     | 1.5,000   | 02.01               |
|                |            | 9       | 1          |            |          |       |        |            |            |         |         |         |             | Score =   | 220                 |

SI Table 6b: Karst Aquifers

| · · · · · · · · · · · · · · · · · · · | · · · · · · · · · · · · · · · · · · · | Nearest    |    | Pop         | ulation | Serve | d by We | Us Within | Distance | Categor | Y       |         |           |               | 1          |
|---------------------------------------|---------------------------------------|------------|----|-------------|---------|-------|---------|-----------|----------|---------|---------|---------|-----------|---------------|------------|
|                                       | 1                                     | Well       | ,  | 11          | 31      | 101   | 301     | 1,001     | 1,001    | 10.001  | 30,001  | 100,001 | 300,001   | 1,000,001     | Population |
|                                       | 1                                     | (use 20    | 10 | lo          | to      | 60    | 10      | 10        | lo       | 10      | 10      | 10      | 10        | to.           |            |
| Distance                              | On a tallon                           | (or karst) | 10 | 30          | 100     | 300   | 1,000   | 3.000     | 10,000   | 30,000  | 100.000 | 300,000 | 1,000,000 | 3,000,000     | Value      |
| trom Site                             | Population                            | 7() XB(31) |    |             |         |       |         |           |          |         | 1       |         |           |               |            |
| ·                                     |                                       | 20         |    | 2           | 6       | 16    | 52      | 163       | 521      | 1.633   | 5,214   | 18,325  | 52,138    | 163,246       |            |
| O to % mile                           |                                       | ]          |    |             |         | 1     |         |           |          |         | 3,233   | 10,121  | 32,324    | 101,212       |            |
| > % to % mile                         | <u> </u>                              | 20         | 1  | 1           | 3       | 10    | 32      | 101       | 323      | 1,012   | 3,233   | 10,121  |           | , , , , , , , |            |
| 7 7 10 10 11                          |                                       |            |    |             | ا . ا   |       | 26      | 82        | 261      | 816     | 2,607   | 8,182   | 26,068    | 81,623        |            |
| > is to 1 mile                        |                                       | 20         | 1  | 1           | ١ ،     | 8     | 40      | • • •     | •••      |         |         |         | ì         |               | }          |
|                                       |                                       | 1 20       | ,  | ١,          | 3       | l a   | 26      | 82        | 261      | 816     | 2,607   | 8,162   | 26,068    | 81,623        |            |
| > 1 to 2 miles                        |                                       | 20         | '  | l '         | 1       | _     | 1       |           |          |         | 1       |         |           |               | 1          |
|                                       |                                       | 20         | 1  | 1           | 3       | 8     | 26      | 82        | 261      | 816     | 2,607   | 8,162   | 26,068    | 81,623        |            |
| > 2 to 3 miles                        |                                       | 1 "        |    | [           | 1       | Į.    | ]       |           |          |         | 1       | 8,162   | 28,068    | 81,623        |            |
| > 3 to 4 miles                        |                                       | 20         | 1  | 1           | 3       | 8     | 26      | 82        | 261      | 816     | 2,607   | 1 3,102 | 1.000     | 1             |            |
|                                       |                                       |            |    | <del></del> |         |       |         |           |          |         |         |         |           | Score =       | 1          |


Nenrest Well =

1

#### GROUND WATER PATHIVAY (concluded)

Data WASTE CHARACTERISTICS Score Troe 8. If any Actual Contamination Targets exist for the ground water pathway, assign the hazardous waste quantity score calculated on page 4, or a score of 100, whichever is 10 GREATER; if no Actual Contamination Targets exist, assign the hazardous waste quantity score calculated on page 4. 100 9. Assign the highest ground water toxicity/mobility value from SI Table 3 or 4. 10. Multiply the ground water toxicity/mobility and waste quantity scores. Assign the Waste Characteristics score from the table below: WC Score Product Product WC Score WC = 10.000 to < 1E+0 ٥ 0 10 >0 to <10 ١ 1E+05 to < 1E+0 18 32 1E+06 to < 1E+0 10 to < 100 2 1E+07 to <1E+0 56 100 to < 1,000 1E+08 or creater 100 1.000 to < 10.000 6

**GROUND WATER PATHWAY SCORE:** 



Parcents

| Sample 10  | Hazardous<br>Substance                                   | Toxicity/<br>Persistence | Toxicity/ Persistence/ Biosccumulation      | Ecolonicity/ Persistence/Eco- blosccumsdation | Reference                    | No sa                                      | amples<br>Supp<br>J cont | taker<br>ort                    | )<br>]<br>3   |
|------------|--|--------------------------|---|---|------------------------------|--|--------------------------|---------------------------------|---------------|
|            |  |                          |   |   |                              | 54   | 1 conf                   | tamij<br>                       | iation<br>i   |
| •          | Highest<br>Values<br>FACE WATER DRINKING W<br>Sample Typ |                          |   | AMINATION                                     |                              | Ро   | pulation Serve           |                                 |               |
| Sample ID  | Hazardous<br>Substance                                   | Concentration (ug/L)     | Benchmerk Gencentration [MCL or MCLG)       | Percent of<br>Benchmerk                       | Cancar Risk Concentration    | Percent of Cancer Risk Concentration       | Reference<br>Dose        | Percent of<br>Reference<br>Dose | í             |
|            |  | <b>↓</b>                 | Highest<br>Percent                          | II.   | Sum of<br>Percents           |  | Sum of<br>Percents       |                                 | J 80 5        |
| Intake ID: | Sample Typ   | )e:                      |   | Level I                                       | Level II                     | Po   | pulation Serve           | d                               | June 30, 1992 |
| Semple ID  | Hezerdous<br>Substance                                   | Concentration<br>(ug/L)  | Benchmark<br>Concentration<br>(MCL or MCLQ) | Percent of<br>Benchmark                       | Cencer Risk<br>Concentration | Percent of<br>Cencer Riek<br>Concentration | Reference<br>Dase        | Percent of Reference Dose       | 992           |
|            |  |                          | Muhasi                                      |   | - Surn el                    |  | Sum of                   |                                 |               |

Percent

Toxicity/

Ecotonicity/

Percents

SI TABLE 7: SURFACE WATER OBSERVED RELEASE SUBSTANCES

NAV 5 1991

| 1. CBSERVED RELEASE: If sampling data or direct observation support a release to surface water, assign a score of 550. Record observed release substances on SI Table 7.  2. NO OBSERVED RELEASE: If sampling data do not support a release to surface water, use the table below to assign a score based on distance to surface water and flood frequency.  Distance to surface water < 2500 feet 500  Distance to surface water > 2500 feet, and  Site in annual or 10-yr floodplain 500  Site in 100-yr floodplain 300  Site outside 500-yr floodplain 100  LR = 100  DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6. | 28<br>No<br>86             |
|---|----------------------------|
| water, assign a score of 550. Record observed release substances on SI Table 7.  2. NO OBSERVED RELEASE: If sampling data do not support a release to surface water, use the table below to assign a score based on distance to surface water and flood frequency.    Distance to surface water < 2500 feet   | <b>88</b>                  |
| Distance to surface water < 2500 feet 500  Distance to surface water > 2500 feet, and 500  Site in annual or 10-yr floodplain 500  Site in 100-yr floodplain 300  Site outside 500-yr floodplain 100  DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.   | <b>88</b>                  |
| Distance to surface water > 2500 feet, and  Site in annual or 10-yr floodplain  Site in 100-yr floodplain  Site in 500-yr floodplain  Site outside 500-yr floodplain  Site outside 500-yr floodplain  LR = 1000  DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit. 5, and 6.   |                            |
| Distance to surface water > 2500 feet, and  Site in annual or 10-yr floodplain  Site in 100-yr floodplain  Site in 500-yr floodplain  Site outside 500-yr floodplain  DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, 3, and 6.  |                            |
| Site in annual or 10-yr floodplain 500 Site in 100-yr floodplain 400 Site in 500-yr floodplain 300 Site outside 500-yr floodplain 100  ER = 100  CRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, 3, and 6.   |                            |
| Site in 500-yr floodplain  Site outside 500-yr floodplain  Site outside 500-yr floodplain  LR = 100  CRINKING WATER THREAT TARGETS  B. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.  |                            |
| Site outside 500-yr floodplain 100  LR = \_\OC\_  DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.   |                            |
| DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.   |                            |
| DRINKING WATER THREAT TARGETS  3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.   |                            |
| 3. Record the water body type, flow (if applicable), and number of people served by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.  | 0-4                        |
| by each drinking-water intake within the target distance limit. If there is no drinking-water intake within the target distance limit, assign 0 to factors 4, 5, and 6.   | b-r                        |
| intake within the target distance limit, assign 0 to factors 4, 5, and 6.   |                            |
|   | 1 he                       |
| Class Papple Served   | 06                         |
| Intake Name Water Body Type Flow People Served  | 85;                        |
| cfs   | 28                         |
| cfs   |                            |
| cfs   |                            |
| Are any intakes part of a blended system? Yes No No If yes, attach a page to show apportionment calculations.   | Pef                        |
|   | Nos                        |
| 4. ACTUAL CONTAMINATION TARGETS: If analytical evidence indicates that any drinking-  | <b>95</b> ;                |
| water intake listed above has been exposed to a hazardous substance from the site, list the   |                            |
| intake name and calculate the factor score based on the intake population from SI Table 8.  | 28                         |
|   |                            |
|   |                            |
| Level I: people x 10 = Total =  |                            |
| 5. POTENTIAL CONTAMINATION TARGETS: Determine the number of people served by  | Sec                        |
| drinking-water intakes that have not been exposed to a hazardous substance from the site.   | NOS<br>25;                 |
| and assign the total population score from SI Table 9.  | 26                         |
| and essign the total population seems.  |                            |
| 6. NEAREST INTAKE: Assign a score of 50 for any Actual Contamination Targets.   | Res                        |
| Action a score of 45 if there are Level II targets but no Level I targets. If no Actual   | No                         |
| Contamination Targets exist, assign the nearest intake score from SI Table 9. If no   | 125                        |
| drinking-water intakes exist, assign 0.   | Dec                        |
| 7. RESOURCES  | 15<br>25<br>26<br>35<br>35 |
| T= C  | )     -                    |
| <del></del>   |                            |
|   |                            |

SURFACE WATER PATHWAY

## There are no Surface water intakes.

#### SI TABLE 9: VALUES FOR POTENTIAL CONTAMINATION SURFACE WATER TARGET POPULATIONS

|   |             | Nearest                       |               |                 | Pop              | ulation            | Served by            | Intakes | Within Flo             | w Catego                | N                        |                            |                              |                   | r                   |
|---|-------------|-------------------------------|---------------|-----------------|------------------|--------------------|----------------------|---------|------------------------|-------------------------|--------------------------|----------------------------|------------------------------|-------------------|---------------------|
| Surlace Water<br>Body Flow<br>(see SI Table 11) | Population  | Intaka<br>(choosa<br>highast) | 1<br>10<br>30 | 31<br>10<br>100 | 101<br>(*<br>300 | 301<br>10<br>1,000 | 1,001<br>to<br>3,000 | 10.000  | 10,001<br>to<br>30,000 | 30,001<br>to<br>100,000 | 100,001<br>ta<br>300,000 | 300,001<br>te<br>1,000,000 | 1,000,001<br>60<br>3,000,000 | 3,000,001<br>(**) | Population<br>Value |
| < 10 cls  |             | 20                            | 2             | 5               | 16               | 52                 | 163                  | 521     | 1,633                  | 5,214                   | 16,325                   | 52,136                     | 163,246                      | 521,359           | -A/                 |
| 0 to 100 cfs                                    |             | 2                             | 1             | 1.              | 2                | 5                  | 16                   | 62      | 163                    | 521                     | 1,633                    | 5,214                      | 16,325                       | 52,136            | 4/                  |
| 100 to 1,000 cls                                |             | 1                             | 0             | ٥               | 1                | 1                  | 2                    | б       | 16                     | 52                      | 163                      | 521                        | 1,633                        | 5,214             | /                   |
| 1,000 to 10,000 cfs                             |             | 0                             | 0             | 0               | ٥                | o                  | 1                    | 1       | 2                      | 5                       | 16                       | 52                         | 163                          | 521               |                     |
| · 10,000 cls or<br>reat Lakes                   | <del></del> | 0                             | 0             | O               | 0                | 0                  | 0                    | ٥       | 1                      | 1                       | 2                        | 5                          | 16                           | 52                |                     |
| nuls Mixing Zons                                |             | 10                            | 1             | 3               | 8                | 26                 | 82                   | 261     | 816                    | 2,607                   | 8,162                    | 26,068                     | 81,623                       | 260,680           |                     |
| Neare   | st Intake = |                               |               |                 |                  |                    |                      |         |                        |                         |                          |                            |                              | Score =           |                     |

SI TABLE 10: SURFACE WATER TYPE / FLOW CHARACTERISTICS WITH DILUTION WEIGHTS FOR POTENTIAL CONTAMINATION SURFACE WATER SENSITIVE ENVIRONMENTS

| Type of Sw   | face Water Body       | Dilution |
|--|-----------------------|----------|
| Water Body Type OR   | Flow                  | Weight   |
| minimal stream   | < 10 cfs              | 1        |
| small to moderate stream   | 10 to 100 cfs         | 0.1      |
| moderate to large stream   | > 100 to 1,000 cla    | N/A      |
| large stream to river  | > 1,000 to 10,000 cfe | N/A      |
| large river  | > 10,000 cfe          | N/A      |
| 3-mile mixing zone of<br>quiet flowing attenns or rivers                       | 10 cfs or greater     | N/A      |
| constal tidal water (haibors,<br>sounds, beys, etc.), ocean,<br>or Great Lakes | N/A                   | N/A      |

1680 g //unc

John Hassall Inc. 8002-064

## No analytical data to Support Contamination to human food Chain

#### SI TABLE 11: HUMAN FOOD CHAIN ACTUAL CONTAMINATION TARGETS

Fishery ID: \_\_\_\_\_ Sample Type: \_\_\_\_ Level II \_\_\_\_


| Sample 10       | Hazardous<br>Substance | Concentration<br>(mg/kg) | Benchmark<br>Concentration<br>(FDAAL)         | Percent of<br>Benchmark | Cancer Risk Concentration | Percent of Cancer Risk Cancentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose |
|-----------------|------------------------|--------------------------|---|-------------------------|---------------------------|--------------------------------------|--------------------|---------------------------------|
|                 |                        |                          | Highest<br>Percent                            |                         | Sum of<br>Percente        |                                      | Sum of<br>Percents |                                 |
|                 | NSITIVE ENVIRONMENT A  |                          | ITAMINATIO                                    |                         |                           | Env                                  | vironment Valu     | e                               |
| Sample ID       | Hezerdaus<br>Substança | Concontration<br>(ug/L)  | Benchmark<br>Concentration<br>(AWQC or AALAC) | Percent of<br>Benchmerk |                           |                                      |                    |                                 |
|                 |                        |                          | Highest<br>Percent                            |                         |                           |                                      |                    |                                 |
| Environment ID: | Sam                    | pla Type:                |   | Lovel I                 | J<br>Lavel II             | Env                                  | rironment Valu     | c                               |
| Sample ID       | Conteminent:           | Concentration<br>(vg/L)  | Bonchmork Concentration [AWGG or AALAG]       | Percent of<br>Benchmerk |                           |                                      |                    |                                 |
| 1               | . ;                    | <u> </u>                 | Highest                                       |                         |                           |                                      |                    |                                 |

## SURFACE WATER PATHWAY (continued) HUMAN FOOD CHAIN THREAT WORKSHEET

17


| 8. Record the water body type and flow (if applicable) for each fishery within the target distance limit. If there is no fishery within the target distance limit, assign a score of 0 at the bottom of this page.    Fishery Name   | core Type | Seferen    |
|--|-----------|------------|
| 9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery                             |           | 95.<br>Nos |
| 9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery                             |           | æ, `       |
| 9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery                             | İ         | 96         |
| 9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery                             |           | 1          |
| 9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery                             | 1         |            |
| <ol> <li>9. ACTUAL CONTAMINATION FISHERIES: If analytical evidence or direct observation indicates that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.</li> <li>10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery</li> </ol> |           | 1 1        |
| that any fishery listed above has been exposed to a hazardous substance from the site, record contaminant information on SI Table 11. Assign a score of 50 if there is a Level I fishery, or 45 for Level II.  10. POTENTIAL CONTAMINATION FISHERIES: If there is a release to a watershed containing fisheries within the target distance limit, but there are no Level I or Level II fisheries, assign a score of 21. If there is no observed release to the watershed, assign a value for potential contamination fisheries from the table below using the LOWEST flow at any fishery   |           |            |
|  | )         |            |
| Lowest Flow Potential Fisheries Score  |           | 11         |
| < 10 cts 21  | ł         |            |
| 10 to 100 cfs 3  | -         |            |
| > 100 cfs, coastal tidal waters,   | 1         | 11         |
| oceans, or Great Lakes   | <b>)</b>  | 14         |

18




### SURFACE WATER PATHWAY (continued) ENVIRONMENTAL THREAT WORKSHEET

े ।ठवा Data **ENVIRONMENTAL THREAT TARGETS** Score Type Refere 11. Record the water body type and flow (if applicable) for each surface water sensitive Det. environment within the target distance limit (see SI Tables 13 and 14). If there is no sensitive environment within the target distance limit, assign a Targets score of O NO at the bottom of the page. Water Body Type Flow Environment Name cfs cfs cts cfs cfs 12. ACTUAL CONTAMINATION SENSITIVE ENVIRONMENTS: If sampling data or direct observation indicates any sensitive environment listed above has been exposed to a hazardous substance from the site, record this information on SI Table 12, and assign a factor score using the environment value from SI Table 13 or 14. Multiplier (10 for Environment Type and Value Level I, 1 for Level III Product (SI Tables 13 or 14) . Environment Name = × 13. POTENTIAL CONTAMINATION SENSITIVE ENVIRONMENTS: A. For Potentially Contaminated Sensitive Environments located on surface water bodies with flows of 100 cfs or less, assign scores as follows: Dilution Weight Environment Type and Value Product ISI Tables 13 or 141 ISI Table 101 Flow × x 0.1 = cts x 0.1 = × cfs × x 0.1 = cfs × x 0.1 = cts x x 0.1 = cfs Sum = B. If any Potentially Contaminated Sensitive Environment is located on coastal tidal waters, ocean, or great lakes, or a surface water body with flow greater than 100 cfs, assign a score of 1.



153




#### SI TABLE 13: SURFACE WATER AND AIR SENSITIVE ENVIRONMENTS VALUES

| Sansitiva Environment  | Assigned Value         |
|--|------------------------|
| Critical habitat for Federally designated andangered or threatened species                                       | 100                    |
| Marine Sanctuary   |                        |
| National Perk  |                        |
| Designated Federal Wilderness Area   |                        |
| Ecologically important areas identified under the Coastal Zone Wilderness Act                                    |                        |
| Sensitive Areas identified under the National Estuary Program or Near Coastal Water Program of the Clean Wat     | ter Act                |
| Critical Areas Identified under the Clean Lakes Program of the Clean Water Act (subareas in lakes or entire sm   | ail lakes)             |
| National Monument (air pathway only)   |                        |
| National Seashore Recreation Area  |                        |
| National Lakeshore Recreation Area   |                        |
| Habitat known to be used by Federally designated or proposed endangered or threatened species                    | 75                     |
| National Preserve  |                        |
| National or State Wildlife Refuge  |                        |
| Unit of Caastal Barner Resources System  |                        |
| Federal land designated for the protection of natural ecosystems   |                        |
| Administratively Proposed Federal Wilderness Area  |                        |
| Spawning areas critical for the maintenance of fish/shellfish species within a river system, bay or estuary      |                        |
| Migratory pathways and feeding areas critical for the maintenance of anacizmous fish species within river rea    | ches                   |
| or areas in lakes or coastal tidal waters in which the fish spend extended periods of time                       |                        |
| Terrestrial areas utilized by large or dense aggregations of vertebrate animas (semi-equatic foragers) for breed | ling                   |
| National over reach designated as recreational   |                        |
| Habitat known to be used by State designated endangered or threatened species                                    | 50                     |
| Habitat known to be used by a species under review as to its Federal endangered or threatened status             |                        |
| Coastal Berrier (partially developed)  |                        |
| Federally designated Scenic or Wild River  |                        |
| State land designated for wildlife or game management  | 25                     |
| State designated Scenic or Wild River  |                        |
| State designated Natural Area  |                        |
| Particular areas, relatively small in size, important to maintenance of unique biotic communities                |                        |
| State designated areas for the protection or maintanance of aquatic life under the Clean Water Act               | 5                      |
| See SI Table 15 (S   | iurlace Weter Pathwey) |
| Wetlands   | or                     |
| Si Table 2   | O (Air Pathway)        |

## SI TABLE 14: SURFACE WATER WETLANDS FRONTAGE VALUES

| Assigned Value |
|----------------|
| 0              |
| 25             |
| 50             |
| 75             |
| 100            |
| 150            |
| 250            |
| 350            |
| 450            |
| 500            |
|                |



#### SURFACE WATER PATHWAY (concluded) WASTE CHARACTERISTICS, THREAT, AND PATHWAY SCORE SUMMARY

21

| WASTE                                  | CHARACTERISTICS  |   |     |                                  |     |                                   | Score    |  |  |  |  |  |
|--|--|---|-----|----------------------------------|-----|-----------------------------------|----------|--|--|--|--|--|
| (p:<br>on<br>8. <b>ii</b>              | 4. A. If any Actual Contamination Target exists for the surface water pathway (pages, 14, 18, or 17), assign the hazardous waste quantity score calculated on page 4, or a score of 100, whichever is GREATER.  8. If there is no Actual Contamination Target for the surface water pathway, assign the hazardous waste quantity score calculated on page 4. |   |     |                                  |     |                                   |          |  |  |  |  |  |
| below                                  | on the highest value from<br>v. Multiply each by the<br>acteristics score for each   | surface water waste to threat using the table Substance | qu. | annty score, and                 | det | e factors listed ermine the waste | WC Score |  |  |  |  |  |
|  | ing Water Threat   | 10,000  | ×   | 10                               | =   | 1×105                             | 18       | :::::::::::::::::::::::::::::::::::::: |  |  |  |  |
| Food (                                 | Chain Threat ity/Persistence/  | 5×104   | x   | 10                               | =   | 5×10 <sup>5</sup>                 | 18       | of<br>1 0003                           |  |  |  |  |
| Ecoto                                  | onmental Threat<br>exicity/Persistence/<br>osccumulation   | 5×106   | x   | 10                               | 2   | 5x107                             | 56       | af<br>1 0000                           |  |  |  |  |
| ــــــــــــــــــــــــــــــــــــــ | Product  | WC Score  |     | Product                          | _   | WC Score                          |          |  |  |  |  |  |
|  | 0<br>>0 to <10   | 0   |     | 1E+06 to < 1E+<br>1E+07 to < 1E+ |     | 32<br>56                          |          |  |  |  |  |  |

STIDEACE WATER DATHWAY THREAT SCORES

2

3

6

10

18

10 to < 100

100 to < 1,000

1,000 to < 10,000

10,000 to <1E+05

1E+05 to <1E+08

| Threat           | Likelihood of<br>Release (LR) Score<br>(from page 13) | Targets (T) Score | Pathway Waste Characteristics (WC) Score (datermined above) | Threat<br>LR x T x W |
|------------------|---|-------------------|---|----------------------|
| Drinking Water   | 10  | Ø                 | 18  | (mezimun             |
| Human Food Chain | 10  | Ø                 | 18  | (mezimur             |
| Environmental :  | 10  | Ø                 | 56  | (mesumu              |

1E+08 to < 1E+0

1E+09 to < 1E+1

1E+10 to <1E+1

1E+11 to <1E+1

1E+12 or greater

SURFACE WATER PATHWAY SCORE

100

180

320

560

1000

(Drinking Water Threat + Human Food Chain Threat + Environmental Threat)





John Hassall 8002-064 June 30, 1992

# No analytical data to Support contamination of residential Soil

| Residence ID: |                        | Level i                  | Lovel II                     |  |                    |                                 | -         | ı           |   |
|---------------|------------------------|--------------------------|------------------------------|--|--------------------|---------------------------------|-----------|-------------|---|
| Semple ID     | Hazardous<br>Substance | Concentration (mg/kg)    | Cancer Risk<br>Concentration | Parcent of<br>Cancer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dase | Reference | <b>.*</b> . | • |
|               | NA                     |                          |                              |  |                    |                                 |           | en<br>T     |   |
|               |                        |                          | Sum of<br>Percents           |  | Sum of<br>Percents |                                 |           | •           | • |
| Residence ID: |                        | Level I                  | Level II                     |  | Population         | n                               | -         |             |   |
| Sample ID     | Hazaidous<br>Substance | Concentration<br>[mg/kg] | Gencer Rick<br>Concentration | Percent of<br>Gencer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose | Reference |             |   |
|               | NA                     |                          |                              |  |                    |                                 |           |             |   |
|               | •                      |                          | Sum of<br>Percents           |  | Sum of<br>Percents |                                 |           |             |   |
| Residence ID: | ·····                  | Level I                  | Level II                     | <del></del>                                | Population         |                                 | -         |             |   |
| Sample ID     | Hazardoua<br>Subatanca | Concentration<br>(mg/kg) | Cancar Risk<br>Concentration | Percent of<br>Cancer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose | Reference |             |   |
|               | NA                     |                          |                              | •  |                    |                                 |           |             |   |
|               |                        |                          | Sum of<br>Percents           |  | Sum of<br>Percents | <del></del>                     |           |             |   |

#### SOIL EXPOSURE PATHWAY WORKSHEET


| LIKELIHO   | OD OF EXPOSURE  |                         |                        | Score    | Type              |
|------------|---|-------------------------|------------------------|----------|-------------------|
| (depth     | VED CONTAMINATION: If evidence into of 2 feet or less), assign a score of 550 sure score of 0 results in a soil exposure.                   | ); otherwise, assign 0. | Note that a likelihood |          | RE 200            |
|            |   |                         | LE                     | 550      |                   |
| RESIDEN    | T POPULATION THREAT TARGET  | S                       |                        |          |                   |
|            | INT POPULATION: Determine the number  |                         |                        |          | pef               |
|            | nding school or daycare on or within 20 to the concentration level on SI Table 1  |                         |                        |          | NO                |
| Calcula    | Level I:  | people x 1              | 0 =                    | ø        | 23                |
| Assign     | INT INDIVIDUAL: Assign a score of 50 a score of 45 if there are Level II targetion exists, assign 0.  |                         |                        | Ø        | 285<br>203<br>203 |
|            | ERS: Assign a score from the table believely facilities with areas of surficial conformation of Workers  O  1 to 100  101 to 1,000  > 1,000 |                         |                        | 5        | Ref.              |
| l '        | STRIAL SENSITIVE ENVIRONMENTS: Userestrial sensitive environment on an ar   | rea of surficial contam | -                      |          | Res<br>No         |
|            | TOTALIST GENERAL ETTAL GIANTEN TYPE   |                         |                        | <b>8</b> | 23                |
| 6. RESOU   |   |                         | Sum                    | - 10     | Rof               |
|            | RCFS  |                         |                        | 1 (7)    | 1, 63             |
| 0. 11.2500 | RCES  |                         |                        | Ø -      | 128               |



24

## SI TABLE 16: SOIL EXPOSURE PATHWAY TERRESTRIAL SENSITIVE ENVIRONMENT VALUES

| Terrestrial Sensitive Environment   | Assigned Value |
|---|----------------|
| Terrestrial critical habitat for Federally designated endangered or threatened species                    | 100            |
| National Park   |                |
| Designated Federal Wilderness Area  |                |
| National Monument   |                |
| Terrestrial habitat known to be used by Federally designated or proposed threatened or endangered species | 75             |
| National Preserve (terrestrial)   |                |
| National or State terrestrial Wildlife Refuge   |                |
| Federal land designated for protection of natural ecosystems  |                |
| Administratively proposed Federal Wilderness Area   |                |
| Terrestrial areas utilized by large or dense aggregations of animals (vertebrate species) for breeding    |                |
| Terrestrial habitat used by State designated endangered or threatened species                             | 50             |
| Terrestrial habitat used by species under review for Federally designated endangered or threatened status |                |
| State lands designated for wildlife or game management  | 25             |
| State designated Natural Areas  |                |
| Particular areas, relatively small in size, important to maintenance of unique biotic communities         |                |





#### SOIL EXPOSURE PATHWAY WORKSHEET (continued)

#### WASTE CHARACTERISTICS

| 7. Assign | 7. Assign the hazardous waste quantity score calculated on page 4. |                  |                                    |                |         |   |  |  |  |  |
|-----------|--|------------------|------------------------------------|----------------|---------|---|--|--|--|--|
| 8. Assign | . Assign the highest toxicity value from SI Table 15.              |                  |                                    |                |         |   |  |  |  |  |
|           | y the toxicity and waste<br>he table below:                        | quantity scores. |                                    | eristics score |         |   |  |  |  |  |
|           | 0  | 0                | 10,000 to < 1E+05                  | 10             | wc - 18 | 3 |  |  |  |  |
|           | >0 to <10<br>10 to <100<br>100 to <1,000                           | 2 3              | 1E+06 to <1E+07<br>1E+07 to <1E+04 | 32<br>56       |         |   |  |  |  |  |
|           | 1,000 to <10.020   | <u> </u>         | 1E+08 or greater                   | 100            |         |   |  |  |  |  |

RESIDENT POPULATION THREAT SCORE: 550x5x18

550x5x18 LEXTXWC = 82,500

0.6

NEARBY POPULATION THREAT SCORE:

Population within one mile: 13336

2.0

SOIL EXPOSURE PATHWAY SCORE:
Resident Population Threat + Nearby Population Threat

2.6

$$\frac{18 \times 5 \times 550}{82,500} + 2.0$$

## NO AIR SAMPLES TAKEN TO SUPPORT CONTRAMINATION OF AIR.

#### SI TABLE 17: AIR PATHWAY OBSERVED RELEASE SUBSTANCES

| 31 IMPLE 17. MI  | TAILLIA C   | DOLITED II.                  | renos                 |   |                         |                              |  |                    |                                 |
|------------------|-------------|------------------------------|-----------------------|---|-------------------------|------------------------------|--|--------------------|---------------------------------|
| Sample ID:       |             | <u>.</u>                     |                       | Lavel I   | Level II                | <del></del> .                | Distance from                              | Source (mi)        |                                 |
| Hazard<br>Substa |             | Concentration<br>(ug/m3)     | Toxicity/<br>Mobility | Benchmark<br>Concentration<br>INAAGS & MESHAPU  | Percent of<br>Benchmerk | Cencer Risk<br>Concentration | Percent of<br>Cencer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose |
|                  | NH          |                              |                       |   |                         |                              |  |                    |                                 |
|                  |             | Highest<br>Toxicity/Mobility |                       | Highest<br>Percent                              |                         | Sum of<br>Percents           |  | Sum of<br>Percents | I .                             |
| Sample ID:       |             | -                            |                       | Level I   | Level II                |                              | Distance from                              | Source (mi)        |                                 |
| Hazərd<br>Substa |             | Concentration<br>(ug/m3)     | Toxicity/<br>Mobility | Benchmark<br>Concentration<br>IMAAOS or MISMAPS | Parcent of<br>Benchmark | Cancer Risk<br>Concentration | Percent of<br>Gancer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dose |
|                  | UA          |                              |                       |   |                         |                              |  |                    |                                 |
|                  |             | Highest<br>Toxicity/Mobility |                       | Highest<br>Percent                              |                         | Sum of<br>Percente           |  | Sum of<br>Percents | •                               |
| Sample ID:       |             | _                            |                       | Level I   | Level II                | <del></del>                  | Distance from                              | Source (mi)        |                                 |
|                  | 1/1         | Concentration<br>(ug/m3)     | Toxicity/<br>Mobility | Benchmark Concentration (NAAOS or NESHAFI)      | Percent of<br>Benchmark | Cancer Risk<br>Concentration | Percent of<br>Gencer Risk<br>Concentration | Reference<br>Dose  | Percent of<br>Reference<br>Dase |
|                  | <i>'\//</i> |                              |                       |   |                         |                              |  |                    |                                 |
|                  | · ·:        | Highest                      |                       | Highest   |                         | Sum of                       |  | Sum of             |                                 |

26

#### AIR PATHWAY WORKSHEET

| LIKELIHOOD OF RELEASE  | Score    | Type | Referen   |
|--|----------|------|-----------|
| <ol> <li>CBSERVED RELEASE: If sampling data or direct observation support a release to air,<br/>assign a score of 550. Record observed release substances on SI Table 17.</li> </ol> |          |      |           |
| <ol><li>NO OBSERVED RELEASE: If sampling data do not support a release to air, assign a<br/>score of 500.</li></ol>  | 500      |      |           |
| IR -   | 500      |      | -         |
| TARGETS  | 1300     |      | i         |
| 3. ACTUAL CONTAMINATION POPULATION: Determine the number of people subject   |          |      |           |
| to exposure from a release of a hazardous substance to the air. Calculate levels of  |          |      | ļ         |
| exposure on SI Table 17.  Level I: people x 10 =   |          |      |           |
| Level II: people x 1 = Tatal =   | 0        |      | <u> </u>  |
| 4. POTENTIAL TARGET POPULATION: Cetermine the number of people not subject   | ł        |      | pes       |
| to exposure from a release of a hazardous substance to the air, and assign the   | 100      |      | 10        |
| total population score from SI Table 18.   | 109      |      |           |
| 5. NEAREST INDIVIDUAL: Assign a score of 50 if there are any Level I targets.  |          |      | ref<br>No |
| Assign a score of 45 if there are Level II targets but no Level I targets. If no Actual Contamination Population exists, assign the Nearest Individual score from SI Table 18.       | 80       |      | ລາ        |
| 6. ACTUAL CONTAMINATION SENSITIVE ENVIRONMENTS: Sum the sensitive environment  | 0.0      |      |           |
| values (Si Table 13) and wetland acreage values (Si Table 19) for environments subject   | Ì        |      |           |
| to exposure from the release of a hazardous substance to the air.  |          |      |           |
| Sensitive Environment Type Value   |          |      |           |
|  |          |      |           |
|  |          |      |           |
| <del></del>  |          |      |           |
| Wetland Acreage Value  |          |      |           |
|  |          |      |           |
|  |          |      |           |
|  |          |      |           |
|  |          |      | ·         |
| Sum =  | D        |      |           |
| 7. POTENTIAL CONTAMINATION SENSITIVE ENVIRONMENTS: Use SI Table 20 to determine the score for sensitive environments not subject to exposure from a release.                         | 0.05     |      | No.       |
| continue and come for sometime announcements has analysed to anti-control foliated.  | 0.635    |      | 2         |
| 8. RESOURCES   | 0        |      |           |
| 7 <b>=</b>   | 129,625  |      |           |
| , –  | -W 1. WW |      | ł         |

i.

Site Name: Date:

#### SI TABLE 18: VALUES FOR POTENTIAL CONTAMINATION AIR TARGET POPULATIONS

|                                    | 1          | Noarest    |     | Population Within Distance Category |          |     |          |            |        |        |         |         |           |             |            |
|------------------------------------|------------|------------|-----|-------------------------------------|----------|-----|----------|------------|--------|--------|---------|---------|-----------|-------------|------------|
| 1                                  | l          | Individual | . 1 | 11                                  | 31       | 101 | 301      | 1,001      | 3,001  | 10,001 | 30,001  | 100,001 | 300,001   | 1,000,001   |            |
| Distance                           |            | (choose    | 10  | 10                                  | to       | (o  | <b>"</b> | lo         | l•     | 10     | to .    | to      | to        | <b>(•</b>   | Population |
| from Sita                          | Population | highest)   | 10  | 30                                  | 100      | 300 | 1,000    | 3,000      | 10.000 | 30,000 | 100,000 | 300,000 | 1.000.000 | J. 000, 000 | Value      |
| On a source                        | 100        | 20.        | 1   | 2                                   | <b>6</b> | 16  | 52       | 163        | 521    | 1,633  | 5,214   | 16,325  | 52,136    | 163,246     | 5          |
| >0 to % nuls                       | 1,339      | 20         | 1   | 1                                   | ٦        | 4   | 13       | <b>(1)</b> | 130    | 408    | 1,303   | 4,081   | 13,034    | 40,811      | 41         |
| > % to % mds                       | 3187       | 2          | ٥   | o                                   | ı        | ,   | 3        | (3)        | 28     | uv     | 202     | 082     | 2,816     | 0,815       | 9          |
| >% to I mile                       | 9,710      | 1          | 0   | o                                   | o        | 1   | ,        | 3          | 0      | 26     | 83      | 261     | 834       | 2,612       | 8_         |
| > 1 to 2 milos                     | 31,323     | ٥          | o   | o                                   | o        | 0   | 1        | 1          | 3      | 8      |         | 83      | 266       | 833         | 27         |
| > 2 to 3 miles                     | 63,425     | ٥          | o   | 0                                   | o        | o   | ,        | ,          | ı      | 4      | (1)     | 38      | 120       | 376         | 12         |
| > 3 to 4 miles                     | 74,891     | o          | o   | 0                                   | 0        | 0   | 0        | 1          | 1      | 2      | (7)     | 23      | 73        | 229         | <u>7</u>   |
| Nearest Individual - 20 Score - 10 |            |            |     |                                     |          |     |          |            |        | 109    |         |         |           |             |            |

Ref. Nos. 311

SI TABLE 19: AIR PATHWAY VALUES FOR WETLAND AREA

| Wetland Area                  | Assigned Value |
|-------------------------------|----------------|
| Less than 1 acre              | 0              |
| 1 to 50 acres                 | 25             |
| Greater than 50 to 100 acres  | 75             |
| Greater than 100 to 150 acres | 125            |
| Greater than 150 to 200 acres | 175            |
| Greater than 200 to 300 acres | 250            |
| Greater than 300 to 400 acres | 350            |
| Grauter than 400 to 500 acres | 450            |
| Greater than 500 acres        | 500            |

SI TABLE 20: DISTANCE WEIGHTS AND CALCULATIONS
FOR AIR PATHWAY POTENTIAL CONTAMINATION SENSITIVE ENVIRONMENTS

| Distance  | Distance<br>Weight |     | Sens | itive Environment Type and Value<br>(from SI Tables 14 and 20) | Product |
|-----------|--------------------|-----|------|--|---------|
| Oneite    | 0.10               | ×   | Ø    |  | Ø       |
|           |                    | ×   |      |  |         |
| 0-1/4 mi  |                    | ×   | 25   | (wetlands)   | 0.625   |
|           | 0.025              | ×   |      |  |         |
|           |                    | ×   |      |  |         |
| 1/4-1/2mi |                    | ×   | Ø    |  |         |
|           | 0.0054             | ×   |      |  |         |
|           |                    | ×   |      |  |         |
|           |                    | ×   |      |  |         |
| K         | ef. A              | lo. | 21   | Total Engleanments Con-  | 0.625   |

John Hassall Inc. 8002-064 June 30, 1992

#### WASTE CHARACTERISTICS

| ha<br>G | any Actual Contamination Ta<br>azardous waste quantity scor<br>REATER; if there are no Actu<br>ssign the hazardous waste qu | e calculated on parallel Contamination 1 | ge 4, or a score of 100,<br>Targets for the air pathw | whichever is    | 10     |   |
|---------|---|--|---|-----------------|--------|---|
|         | ssign the highest air toxicity/   |  |   |                 | 10     |   |
|         | Aultiply the air pathway toxici   |  | iste quantity scores. As                              | isign the Waste |        |   |
|         | Aultiply the air pathway toxici Characteristics score from the Product  |  | Product   | WC Score        |        | • |
|         | Characteristics score from the  | table below:                             |   |                 | wc = 4 | 3 |
|         | Product   | table below:                             | Product   | WC Score        | wc = 3 | 3 |
|         | Product 0   | table below:                             | Product<br>10,000 to < 1E+0                           | 10<br>18        | wc = 3 | 3 |
|         | Product  O  > 0 to < 10   | table below:                             | Product 10,000 to < 1E+0 1E+05 to < 1E+0              | 10<br>18<br>32  | wc - 3 | 3 |

AIR PATHWAY SCORE

82,500

2.357

500 x 3 x 129.625